

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/304661106>

Variation of Abnormal Hemoglobins Concentrated in Durg, Chhattisgarh: A Brief Note Based on Cross-Sectional Study

Article · January 2016

DOI: 10.4172/2327-5790.1000135

CITATIONS

0

READS

63

2 authors, including:



[Devendra Lingojwar](#)

Albert Einstein College of Medicine

101 PUBLICATIONS 93 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Genomics for Ayurveda [View project](#)



Gene Sequence [View project](#)



Case Report

Variation of Abnormal Hemoglobins Concentrated in Durg, Chhattisgarh: A Brief Note Based on Cross-Sectional Study

Devendra Lingojwar^{1,2,4*}, Pramod Gupta², Savita Bhutoria⁴, Sarita Lingojwar^{1,2}, Nikhil Mishra³ and Anil Kumar³

Abstract

Prevalence of different abnormal hemoglobins (Hb) in Indian tribal and nontribal population groups is well established. Sick cell hemoglobin (HbS) is mostly concentrated in Central and South Indian states, whereas HbD and Hb E is mostly found in North-North-West and North-North-East states respectively. HbJ, an alpha globin gene variant, is earlier reported in North India whereas; its presence in the tribal Chhattisgarh state is not well understood. HbE, a beta-globin gene variant was earlier reported in North Eastern states of India. Prevalence of both these abnormal hemoglobins in the Central India specifically in Durg, Chattisgarh is incompletely understood. In this study attempts were made to analyze the presence of abnormal hemoglobins during screening for sickle cell anemia. Briefly, blood samples (N=44) were analyzed for sickle cell anemia screening and confirmatory tests by solubility tests and cellulose acetate membrane electrophoresis at alkaline pH. Two samples showed an abnormal pattern of separation on cellulose acetate membrane other than HbS. Out of total 44 tested samples, five were sickle cell carriers (HbAS), one was heterozygous HbAJ and the another one was homozygous for HbE while remaining other were normal genotypes i.e. HbAA. In brief, a case of homozygous Hemoglobin E from Kurmi caste of other backward community (OBC) and a different heterozygote pattern, i.e. Hb AJ from Brahmin community is reported from Durg, Chattisgarh, Central India. This study provides possible indication of variation of different abnormal hemoglobins, other than HbS, present in the tribal state of Chhattisgarh.

Keywords

Abnormal hemoglobins; Hemoglobin E; Hemoglobin J; Sickle cell carriers; Indian tribe; Chhattisgarh; Endemic malaria

Case Study

The origin of hemoglobinopathies and prevalence of endemic malaria are inter-linked. Natural protection from malaria pathogenesis after infection might be the most likely and highly accepted hypothesis of the origin of different alleles of normal adult hemoglobin in different population groups throughout the world,

more specifically in the tropical countries where malaria is endemic. Mutation in some of the of genes encoding hemoglobin, red cell enzymes and membrane proteins are being extensively studied with reference to protection from *Plasmodium falciparum* [1]. Hemoglobin S is very common in tribal Indian states upto 35% carrier prevalence in states of Maharashtra [2].

HbE is one of the world's most common and important mutation. It results in a heterogeneous group of disorders whose phenotype range from asymptomatic to severe. HbE trait and Hb EE are mild disorders [3]. The Hb E β^{26} (Glu to Lys) is concentrated in parts of South East Asia where malaria is endemic and HbE carrier status has been shown to confer some protection against *P. falciparum* malaria [4]. Pathogenesis study suggested that patients who co-inherit a mild β -thalassaemia allele with Hb E may have disease on the mild end of the spectrum while those who co-inherited severe β^+ or β^0 thalassaemia alleles might be more severely affected [5]. First reported in Assam with 23% carrier prevalence, HbE is widely distributed in North Eastern states of India with high prevalence amongst 46.4 % in Ahoms of Assam i.e. one of the highest for any abnormal hemoglobin reported from any population in the world. Interestingly, only 1% in Mizoram, 3-33% in West Bengal, while it is almost non-existing in South India [6]. Chhattisgarh is a tribal state and well known for the high prevalence of *P. falciparum* malaria. Presence of abnormal hemoglobin i.e. HbS which is other than normal adult hemoglobin (Hb A), is earlier reported and its prevalence pattern is being studied by different groups [7-10]. Prevalence of this abnormal hemoglobin in Central India specifically in Durg, Chhattisgarh is incompletely understood. Attempts were made to analyze urban samples during sickle cell camp at Durg for finding Hb E in the same population group.

There are more than 50 hemoglobin J variants described in the literature [11]. They all have an electrophoretic mobility "faster" than HbA towards anode in common. All are classified under "variants of the alpha or beta chains" (single or multiple base changes) or "hemoglobins with more than one amino acid substitution in the alpha chain." Hemoglobin J, depending on its type, have different characteristics and functions. Indian variant of Hemoglobin J was reported earlier in North Indian region [12]. Previously Hemoglobin J has been noted by many researchers in various countries. The case of HbJ-Rajappen was reported by Hyde et. al. and later by Henthorn et. Al. in their results of a 10-year program in an English Health region [13,14]. HbJ Baltimore was first described in 1963 in an African-American family. Since then, several cases have been reported in distinct racial groups and also incidentally during the study of other entities, such as thalassaemia [15]. More recently, the increasingly frequent determination of HbA1c in diabetic persons has contributed to the appearance of cases of Hb J-Baltimore associated with anomalous HbA1c values [16,17].

In brief, separation of hemoglobin composition was carried out by cellulose acetate membrane electrophoresis method after solubility tests. Briefly, peripheral blood samples (N=44) after red cell washing and solubility testing for screening HbS were processed for hemolysate preparation by osmotic shock method. Pure hemoglobin from hemolysate was subjected to cellulose acetate membrane electrophoresis in Tris-Glycine buffer (pH 8.6). After 45 minutes, resultant hemoglobin pattern was stained by Ponceau S red

*Corresponding author: Dr. Devendra Lingojwar, Hematology Division, Department of Medicine, Albert Einstein College of Medicine, Bronx NY, USA 10461, Tel: +1 7169512776; E-mail: dplingojwar@gmail.com

Received: April 25, 2016 Accepted: May 19, 2016 Published: May 19, 2016

dye followed by destaining by 5% acetic acid. Appropriate known controls i.e. sickle cell carrier samples were applied every time new samples loaded on cellulose acetate membrane for electrophoretic separation of samples.

Out of 44 samples as shown in Table 1, five samples showed sickle cell carrier status with one fast migrating (HbA) and one slow migrating band (HbS). Two samples i.e. #17 and #27 (Figures 1 and 2) showed mobility pattern different than HbS. Based on mobility pattern, both samples were compared with normally fast migrating HbA and slow migrating HbS. Homozygous Hemoglobin E formed

Table 1: All three abnormal hemoglobin's along with normal pattern (Hb A+A) shown, around 15% abnormal hemoglobin's (Hb A+S, Hb E+E and Hb A+J) were found in the study.

Gender	Mean age in years	Hemoglobin electrophoresis on cellulose acetate membrane in alkaline conditions			
		HbA+A	HbA+S	HbA+J	HbE+E
Male (n=09)	26.77	09	0	0	0
Female (n=35)	22.08	28	5	1	1
Total (n=44)	23.04	37	5	1	1
Percent prevalence		84.09	11.36	2.27	2.27




Figure 1: The family tree was shown. The man pointed by an arrow is the proband. The man A had a renal disease but the detail is unclear. The woman B suffers from recurrent self-limiting febrile episodes and arthralgia but the diagnosis is not made. The man C is on hemodialysis. The woman D had a renal transplant.




Figure 2: Hemoglobin electrophoresis for sample # 17 and 27. Known A+S sample was applied in between #17 and #27 as appropriate control for referring position of HbA, and HbS on the gel. One homozygous sickle cell disease sample (HbS+S) available in the lab from earlier studies* was also applied below sample #17 as a second positive control for examining position of HbE+E and first positive control HbA+S.

a single band and was very slow during electrophoresis and resolved before HbS whereas, heterozygous HbAJ formed one band with similar mobility pattern like HbA for its majority of fractions and second fast migrating band faster than HbA. Separation pattern of sample no. 17 was different from that of HbA and HbS. After comparison with these abnormal hemoglobins, known mobility of Hb E from review of literature and repeated application during cellulose acetate membrane electrophoresis, it is confirmed that this is Hb E.

In brief, in the present study, a case of homozygous HbE from Kurmi caste of OBC community is found in one 21 year old normal healthy female in Durg Chhattisgarh, Central India. Along with this one abnormal heterozygous case of abnormal hemoglobin variant, HbAJ was also found in one 20 year old healthy female from the Brahmin community without any associated clinical manifestations. As this study was based on very small sample size, the presence of different hemoglobin variants such as HbE and HbJ in single study provides possible indication of variation of different abnormal hemoglobins, other than HbS, present in the tribal state of Chhattisgarh. This initial finding of the presence of abnormal hemoglobin variant in the tribal Chhattisgarh state may prove a significant step for further careful observation of routine sickle cell anemia screening.

Acknowledgement

Authors are thankful to the staff of ATG LAB for their kind help and support for lab studies and Department of Biotechnology, Government V.Y.T. PG. Autonomous College, Durg for their local hospitality and arrangement during the diagnostic field work.

Author contributions

Devendra Lingojwar (DL) contributed to the conception and design of the project, prepared the manuscript and revised it critically for intellectual content. PG, SL and NM has acquired, analyzed and interpreted the data, contributed to data collection and analysis. DL, SB and AK has contributed in image analysis and manuscript preparation. All authors approved the final version of the article.

Grant information

This project was conducted as a part of an intramural project of ATG LAB and no specific grant was used for this project from any funding agency.

References

- Allison AC (1954) The distribution of the sickle-cell trait in East Africa and elsewhere, and its apparent relationship to the incidence of subtertian malaria. *Trans R Soc Trop Med Hyg* 48: 312-318.
- Kate S, Lingojwar D (2002) Epidemiology of sickle cell disorder in the state of Maharashtra. *Int J Hum Genet* 2: 161-167.
- Vichinsky E (2007) Hemoglobin e syndromes. *Hematology Am Soc Hematol Educ Program*.
- Ohashi J, Naka I, Patarapotikul J, Hananantachai H, Brittenham G, et al. (2004) Extended linkage disequilibrium surrounding the hemoglobin E variant due to malarial selection. *Am J Hum Genet* 74: 1198-1208.
- Winichagoon P, Thonglairoam V, Fucharoen S, Wilairat P, Fukumaki Y, et al. (1993) Severity differences in beta-thalassaemia/haemoglobin E syndromes: implication of genetic factors. *Br J Haematol* 83: 633-639.
- Agarwal MB (2005) The burden of haemoglobinopathies in India--time to wake up? *J Assoc Physicians India* 53: 1017-1018.
- Panigrahi S, Patra PK, Khodiar PK (2012) Neonatal screening of sickle cell anemia: a preliminary report. *Indian J Pediatr* 79: 747-750.
- Balgir RS (2012) Community expansion and gene geography of sickle cell trait and G6PD deficiency, and natural selection against malaria: experience from tribal land of India. *Cardiovasc Hematol Agents Med Chem* 10: 3-13.
- Lingojwar D, Kate S, Gore M, Basu A (2004) Prevalence of human parvovirus B19 in some tribal population groups from India. *International symposium on*

Emerging Viral Infections: New Frontiers & Challenges.

- Bhagat S, Patra PK, Thakur AS (2013) Fetal Haemoglobin and β^2 -globin Gene Cluster Haplotypes among Sickle Cell Patients in Chhattisgarh. *J Clin Diagn Res* 7: 269-272.
- Patrinos GP, Giardine B, Riemer C, Miller W, Chui DH, et al. (2004) Improvements in the HbVar database of human hemoglobin variants and thalassaemia mutations for population and sequence variation studies. *Nucleic Acids Res* 32: 537-541.
- Srinivas U, Mahapatra M, Pati HP (2007) Hb J Meerut, a fast-moving hemoglobin: a study of seven cases from India and a review of literature. *Am J Hematol* 82: 666-667.
- Hyde RD, Kinderlerer JL, Lehmann H, Hall MD (1971) Haemoglobin J Rajappen; 90 (FG2) Lys leads to Thr. *Biochim Biophys Acta* 243: 515-519.
- Almeida AM, Henthorn JS, Davies SC (2001) Neonatal screening for haemoglobinopathies: the results of a 10-year programme in an English Health Region. *Br J Haematol* 112: 32-35.
- Arribalzaga K, Ricard MP, Carreño DL, Sanchez J, Gonzalez A, et al. (1996) Hb J-Baltimore [β 16(A13)Gly \rightarrow Asp] associated with β (+)-thalassaemia in a Spanish family. *Hemoglobin* 20: 79-84.
- Vandenesch F, Baklouti F, Francina A, Vianey-Liaud C, Bertrand A, et al. (1987) Hemoglobin J-Baltimore (β 16(A13)Gly \rightarrow Asp): interference with the assay of HbA1c. *Clin Chim Acta* 168: 121-128.
- Barakat O, Krishnan ST, Dhatariya K (2008) Falsely low HbA1c value due to a rare variant of hemoglobin J-Baltimore. *Prim Care Diabetes* 2: 155-157.

Author Affiliation

[Top](#)

¹ATG LAB, Biotechnology Research Laboratory, Ganesh Nagar, Pimple Nilakh, Pune, India

²RESEARCH (Regional Society for Education and Research in Community Health), Pune, India

³Department of Biotechnology, Govt. V.Y.T. PG. Autonomous College, G.E. Road, Durg Chhattisgarh, India

⁴Hematology Division, Department of Medicine, Albert Einstein College of Medicine, Bronx NY, USA

Submit your next manuscript and get advantages of SciTechnol submissions

- ❖ 50 Journals
- ❖ 21 Day rapid review process
- ❖ 1000 Editorial team
- ❖ 2 Million readers
- ❖ Publication immediately after acceptance
- ❖ Quality and quick editorial, review processing

Submit your next manuscript at • www.scitechnol.com/submission